



Addition Reactions

- Occurs on double bonds and triple bonds




Mechanism of hydrogenation



Hydrogenation examples

Example 1: 1,2-dimethylcyclobutene



1,2-dimethylcyclobutene

cis-1,2-dimethylcyclobutane


The hydrogenation can occur from the top or the bottom, which in this case produces the same product (*cis* isomer of 1,2-dimethylcyclobutane). The starting material is achiral, and the product is a **meso compound** (two stereogenic centers, but a plane of symmetry)

Example 2: Fats





Glycerol

Basic components are used to make fatty acids.




mono-unsaturated fat (oil)



Example 3: Pentene

Energy is released in each of these reactions, the energy released implies stabilization caused from transforming the starting material into the product


Energy diagram



$$\Delta G \text{ trans-isomer} < \Delta G \text{ cis-isomer} < \Delta G \text{ 1-pentene isomer}$$


Alkenes with more substituents are more stable. Carbons in a double bond have a δ^+ (electron-deficient), this is stabilized by the **electron donating effects** of alkyl groups. Hydrogens are less electron-donating and so less substituted alkenes are less stable.



Cis alkenes are less stable than trans alkenes as they have methyl groups facing the same direction, which causes unfavorable steric interactions.



Hydrogenation:

Limonene




**Example 4: Steroids**

The deuteriums add to the bottom because of the steric hindrance of the nearby methyl group.


$^2\text{H} = \text{D} = \text{deuterium}$

Halogenation: Addition of halogens across a double bond




Note: Iodine (I_2) does not react with most alkenes (i.e., won't add across the double bond).

Example: 2-Methylpropene



Example: 2-Methylpropene




- The electrons in the π -bond attack the partially positive (δ^+) I


- Cl⁻ will attack the more substituted C due to there being a more carbocation character (more substituted end can stabilize the positive charge better due to stabilizing effect of alkyl groups – **inductive effect**)

Markovnikov's Rule: In an addition reaction, the positive end of an A–B system (e.g. I–Cl) adds to the least substituted end of the double bond to make the more stable carbocation.

Example 1: Cyclopentene





Mechanism:




The first step can happen from the top or bottom, and the S_N2 step happens from the opposite side each time.



Example 2: 1,2-dimethylcyclopentene

**Mechanism:**

- Halogen addition to alkene is very fast at -78°C
- Attack always comes from the backside – *trans* product
- When the bromonium ion forms, the bond between the Br and C atom can stretch, giving more carbocation character to the C.


Example: *trans*-2-butene

- Stereochemistry of starting material determines the stereochemistry of product


Mechanisms:#1) Addition of an alcohol functional group (in H₂O)**Example: Cyclohexene**

- In the presence of high concentration of H₂O (55.5 M), H₂O competes with the halide (X⁻) as a nucleophile and reacts with the bromonium intermediate to form **halohydrin** – 1,2-halo alcohols.
- The same is true when using alcohols (ROH) as solvents instead of water.


#2) Addition of an ether functional group (in CH₃OH - methanol)




Summary:







Will add in Markovnikov fashion




Example: 1,2-dimethylcyclobutene




Example: 1-methylcyclohex-1-ene

**Example: Cyclobutene**

Example: Ethylene (intramolecular)

- NaOH – good nucleophile and can attack a primary alkyl halide (intermolecular)
- NaOH – can act as a base and deprotonate the -OH group. The deprotonated -OH group would then attack the primary alkyl chloride to form an ethylene oxide via an Sn₂ reaction (**intramolecular reaction** – within one molecule)
- **Intramolecular reaction is faster than intermolecular reactions**
- OH bonds break fairly easily so proton transfer reactions happen at a faster rate

Example 10: Intramolecular

- Although iodine cannot form diiodide (I₂ cannot add to double bonds), it can form the iodonium ion.
- In the above example, the intramolecular reaction (meaning within the same molecule) occurs much much **FASTER** than the intermolecular reaction (between two or more molecules). This means that the -OH group will attack the iodonium ion much faster than the I⁻ group because it is an intramolecular reaction.
- Intramolecular reaction almost always beats intermolecular reactions.
- 5-membered rings form much faster than 6-membered rings.